

L7 13 L3(5A)L6  
1. 5,643,747, Jul. 1, 1997, Genes for the export of pertussis  
holotoxin; Steven M. Baker, et al., 435/69.1, 252.3, 252.33,  
320.1; 536/23.7 :IMAGE AVAILABLE:

US PAT NO: 5,643,747 :IMAGE AVAILABLE: L7: 1  
of 13  
DETDSC:DETD(31)  
Identification of B. \*\*pertussis\*\* \*\*toxin\*\* secretion  
\*\*mutants\*\*

L12  
1. 5,681,570, Oct. 28, 1997, Immunogenic conjugate molecules;  
Yan-ping Yang, et al., 424/197.11, 203.1, 234.1, 244.1, 256.1;  
514/54; 536/123.1 :IMAGE AVAILABLE:

2. 5,679,784, Oct. 21, 1997, Recombinant mutants for inducing  
specific immune responses; Daniel Iadant, et al., 536/23.7;  
424/184.1, 240.1; 435/69.3, 252.3, 320.1 :IMAGE AVAILABLE:  
3. 5,667,787, Sep. 16, 1997, Purification of a pertussis  
outer membrane protein; Gail Jackson, et al., 424/253.1,  
254.1; 530/412, 414, 416, 417, 418 :IMAGE AVAILABLE:

4. 5,648,080, Jul. 15, 1997, Antigenic preparations and  
isolation of such preparations; Pavel Novotny, 424/254.1,  
184.1, 240.1, 253.1; 530/350, 388.2 :IMAGE AVAILABLE:

5. 5,643,753, Jul. 1, 1997, Use of autologous promoters to  
express gene products in bordetella; Sheena Loosmore, et al.,  
435/69.3; 424/93.4, 235.1, 240.1, 253.1, 254.1; 435/91.1,  
91.4, 91.41, 172.3, 252.3, 320.1; 536/23.7; 935/10, 12, 29,  
41, 56, 72 :IMAGE AVAILABLE:

US PAT NO: 5,681,570 :IMAGE AVAILABLE: L12: 1  
of 45  
DETDSC: DETD(102)  
6. Schneerson, R., L. Levi, J. B. Robibin, D. M. Bryla, G.  
Schiffman, and T. Lagergard. 1992. Synthesis of a conjugate  
\*\*vaccine\*\* composed of pneumococcus type 14 capsular  
polysaccharide bound to \*\*pertussis\*\* \*\*toxin\*\*. Infect.  
Immun. 60:3528-3532.

US PAT NO: 5,679,784 :IMAGE AVAILABLE: L12: 2  
of 45  
DETDSC: DETD(34)  
B. \*\*pertussis\*\* AC \*\*toxin\*\* is used to present epitopes of  
vaccinal importance to the immune system. This recombinant  
toxin may be used as a component of the \*\*vaccine\*\*, either

Welcome to MESSENGER (APS Text) at USPTO

The USPTO production files are current through:

04 NOV 1997 for U.S. Patent Text Data.

04 NOV 1997 for U.S. Current Classification data.

04 NOV 1997 for U.S. Patent Image Data.

FILE 'USPAT' ENTERED AT 15:23:33 ON 07 NOV 1997

U. S. P A T E N T T E X T F I L E

W E L C O M E T O T H E

OR SUBSTITUT? OR MUTAN? OR MUTAGEN?

REPLAC?

(W)9 OR ARG9 OR ARGinine9

1 L8(P)L1

1 L3 AND L8

4149 VACCINE

45 L11(P)L3

1667 (514/12 OR 424/240.1 OR 424/190.1 OR

424/94.5)/CCLS

44 L3 AND L13

1854 (TOXIN/CLM) OR HOLOTOXIN/CLM OR S1/CLM

29 L14 AND L15

alone or in the presence of other antigenic preparation(s). It can be used in toxic form or in detoxified. . .

US PAT NO: 5,667,787 :IMAGE AVAILABLE: L12: 3  
of 45

SUMMARY: BSUM(5)

This . . . hampered by a lack of information on the mechanism of pathogenesis of B. pertussis. Many virulence associated factors, such as \*\*pertussis\*\* \*\*toxin\*\* (PT), also known as lymphocytosis promoting factor (LPF), filamentous haemagglutinin (FHA), adenylate cyclase, lipopolysaccharide, agglutinogens and other outer membrane proteins have been suggested for inclusion in an "acellular" \*\*vaccine\*\*, which is less defined than the component \*\*vaccine\*\*. Much of the work on acellular vaccines has concentrated on a PT-based \*\*vaccine\*\*. Results of a recent clinical trial have indicated that a \*\*vaccine\*\* consisting entirely of PT-toxoid only partially protected children from the infection. A PT/FHA combination showed slightly higher efficacy but this was still lower than .that obtained for the whole-cell \*\*vaccine\*\*.

#### CLAIMS:

3. The \*\*vaccine\*\* claimed in claim 2, wherein said at least one other purified immunoprotective pertussis antigen is selected from the group consisting of purified \*\*pertussis\*\* \*\*toxin\*\* (PT), purified filamentous haemagglutinin (FHA) and purified agglutinogens.

9. The method of claim 8 wherein the \*\*vaccine\*\* further comprises, as at least one additional component thereof, at least one other purified antigen of Bordetella pertussis selected from the group consisting of purified \*\*pertussis\*\* \*\*toxin\*\* (PT), purified filamentous haemagglutinin (FHA) and purified agglutinogens.

10. A component \*\*vaccine\*\* against disease caused by infection by Bordetella pertussis, which comprises: as a first component, purified pertactin having no detectable adenylate cyclase activity, as an additional component, at least one other purified B. pertussis antigen selected from the group consisting of \*\*pertussis\*\* \*\*toxin\*\* (PT), filamentous haemagglutinin (FHA) and agglutinogens; and a pharmacologically acceptable carrier therefor.

US PAT NO: 5,648,080 :IMAGE AVAILABLE: L12: 4  
of 45

SUMMARY: BSUM(4)

The search for a safer, effectively acellular B. pertussis \*\*vaccine\*\* has been hampered in the past by the paucity of information regarding the identity and mechanisms of action of

the. . . J. Am. Med. Soc., 248 (1) 22-23). Examples of antigens that have been suggested for investigation include lymphocytosis promoting factor (\*\*pertussis\*\* \*\*toxin\*\*/LPF) filamentous haemagglutinin (FHA), lipopolysaccharide (LPS), agglutinogens, dermonecrotic toxin (DNT), heat labile and heat stable toxins, polymorphonuclear leukocyte inhibitor factor, adenylate cyclase and other surface components (Pertussis \*\*Vaccine\*\* Workshop, Feb. 11, 1982, Bureau of Biologics, U.S.A.). Other proposed candidate antigens for investigation include tracheal cytotoxin and various outer. . .

BSUM(7)

Much of the work carried out on acellular pertussis vaccines is concentrated on the possibility of basing such a \*\*vaccine\*\* on LPF. However, it is believed that most (if not all) of the adverse effects hitherto observed to be associated with \*\*pertussis\*\* vaccination are related to the \*\*toxin\*\*. In combination with tetanus or diphtheria toxoid and LPS, it is able to induce experimental encephalopathy in susceptible mice (L. . .

US PAT NO: 5,643,753 :IMAGE AVAILABLE: L12: 5  
of 45

SUMMARY: BSUM(4)

The . . . whooping cough with the former generating more severe symptoms. The disease, or vaccination against the disease (using an inactivated whole-cell \*\*vaccine\*\*), elicits antibodies against several antigens, typically \*\*pertussis\*\* \*\*toxin\*\* (PT), filamentous haemagglutinin (FHA), agglutinogens or fimbriae and the 69 kDa outer membrane protein or pertactin. These proteins represent the major immunogens that may be included, individually or in combination, in any \*\*vaccine\*\* used to protect against the disease, whether it be the inactivated whole-cell \*\*vaccine\*\* or a defined component \*\*vaccine\*\*. Therefore, the efficient expression of these antigens from the \*\*vaccine\*\* strain is crucial.

BSUM(7)

The . . . vaccines has resulted in a massive reduction in the incidence of whooping cough since their introduction in the 1950s. These \*\*vaccine\*\* preparations are efficacious but have been known for many years to be reactogenic and to be associated with local and. . . are at various stages of clinical assessment in other countries. Defined pertussis vaccines consist of several combinations of the B.

\*\*pertussis\*\*-specific antigens \*\*pertussis\*\* \*\*toxin\*\* (PT), filamentous haemagglutinin (FHA), the 69 kDa outer membrane protein (pertactin) and fimbrial agglutinogens. Replacing whole-cell whooping cough vaccines with the defined acellular

preparations has resulted in a substantial increase in the complexity and cost of \*\*vaccine\*\* manufacture. A major portion of these increased costs is due to the relatively low levels of PT and pertactin produced. . .

L16

1. 5,643,753, Jul. 1, 1997, Use of autologous promoters to express gene products in bordetella; Sheena Loosmore, et al., 435/69.3; 424/93.4, 235.1, \*\*240.1\*\*, 253.1, 254.1; 435/91.1, 91.4, 91.41, 172.3, 252.3, 320.1; 536/23.7; 935/10, 12, 29, 41, 56, 72 :IMAGE AVAILABLE:

US PAT NO: 5,643,753 :IMAGE AVAILABLE: L16: 1 of 29  
US-CL-CURRENT: 435/69.3; 424/93.4, 235.1, \*\*240.1\*\*, 253.1, 254.1; 435/91.1, 91.4, 91.41, 172.3, 252.3, 320.1; 536/23.7; 935/10, 12, 29, 41, 56, 72

SUMMARY: BSUM(4)

The . . . severe symptoms. The disease, or vaccination against the disease (using an inactivated whole-cell vaccine), elicits antibodies against several antigens, typically \*\*pertussis\*\* \*\*toxin\*\* (PT), filamentous haemagglutinin (FHA), agglutinogens or fimbriae and the 69 kDa outer membrane protein or pertactin. These proteins represent the. . .

BSUM(6)

Along with other genes, the \*\*pertussis\*\* \*\*toxin\*\* operon (TOX), the filamentous haemagglutinin operon (FHA), and the pertactin gene (PRN) are all positively regulated by the Bordetella virulence.

BSUM(7)

The . . . are at various stages of clinical assessment in other countries. Defined pertussis vaccines consist of several combinations of the B. \*\*pertussis\*\* specific antigens \*\*pertussis\*\* \*\*toxin\*\* (PT), filamentous haemagglutinin (FHA), the 69 kDa outer membrane protein (pertactin) and fimbrial agglutinogens. Replacing whole-cell whooping cough vaccines with. . .

DEDESC: DETD(18)

A pUC-based plasmid (S-3680-10) was constructed which contains the FHA promoter directing the expression of the native \*\*pertussis\*\* \*\*toxin\*\* structural gene and surrounded by the TOX flanking regions (FIG. 1). The FHA promoter is an EcoR I/Hinf I fragment. . .

DETD (64)

TABLE 1

Production of \*\*pertussis\*\* \*\*toxin\*\* (PT), filamentous haemagglutinin (FHA) and pertactin, from Bordetella pertussis strains that contain hybrid prn alleles at the prn locus. The. . .

CLAIMS:

What . . . a Bordetella gene fused at the ATG start codon to an autologous Bordetella promoter, wherein: (a) said promoter is the \*\*pertussis\*\* \*\*toxin\*\* operon (TOXp) promoter and the Bordetella gene is selected from the group consisting of the filamentous haemagglutinin operon (FHA) and. . . said promoter is the filamentous haemagglutinin operon (FHAp) promoter and the Bordetella gene is selected from the group consisting of \*\*pertussis\*\* \*\*toxin\*\* operon (TOX) and pertactin (PRN) gene.

3. . . gene fused at an ATG start codon to its own native Bordetella promoter, wherein: (a) said autologous promoter is the \*\*pertussis\*\* \*\*toxin\*\* operon (TOXp) promoter and the Bordetella gene is selected from the group consisting of the filamentous haemagglutinin operon (FHA) and. . . promoter is the filamentous haemagglutinin operon (FHA) promoter and the Bordetella gene is selected from the group consisting of the \*\*pertussis\*\* \*\*toxin\*\* operon (TOX) and pertactin (PRN) gene.

12. . . a Bordetella gene fused at the ATG start codon to an autologous Bordetella promoter, wherein: (1) said promoter is the \*\*pertussis\*\* \*\*toxin\*\* operon (TOXp) promoter and the Bordetella gene is selected from the group consisting of the filamentous haemagglutinin operon (FHA) and. . . said promoter is the filamentous haemagglutinin operon (FHAp) promoter and the Bordetella gene is selected from the group consisting of \*\*pertussis\*\* \*\*toxin\*\* operon (TOX) and pertactin (PRN) gene; (b) introducing said hybrid gene into a Bordetella strain to form a viable transformed. . .

08/483,326

2

Trying 01083...Open

PLEASE ENTER HOST PORT ID:  
PLEASE ENTER HOST PORT ID:X  
LOGINID:d184geb  
PASSWORD:

TERMINAL (ENTER 1, 2, 3, 4, OR ?): 3

including any warranty of marketability of fitness for a particular purpose; nor assumes any legal liability or responsibility for any party's use, or the results of such, of the data.

Help Desk --> 703-305-9000

The Help Desk is staffed for APS support 7 days/week.

Monday through Friday: 6:30am - 9:00pm

Saturday, Sunday, Holidays: 8:30am - 5:00 pm

The Help Desk staff at this number will handle all APS related questions.

[illegible]

The APS is available:

6:30am - 9:00pm Monday through Friday

7:30am - 5:00pm Saturday, Sunday, Holidays

APS is unavailable Thanksgiving Day, Christmas Day, and New Year's Day.

\* \* \* \* \*

FILE 'USPAT' ENTERED AT 13:58:18 ON 04 MAY 1998

\* \* \* \* \*

FILE 'USPAT' ENTERED AT 13:58:18 ON 04 MAY 1998

\*\*\*\*\*

                  welcome to MESSENGER (APS Text) at USPTO

\*\*\*\*\*

The USPTO production files are current through:

APR 28 1998 for U.S. Patent Text Data.

APR 28 1998 for U.S. Current Classification data.

APR 28 1998 for U.S. Patent Image Data.

\*\*\*\*\*

\* \* \* \* \*

\* \* \* \* \*  
 More U.S. patent data is now available on APS. The new  
 USOCR file contains patents issued in 1970, plus some  
 patents that were missing from the USPAT file. See the  
 Patents News Folder under the Public Folders in e-mail for  
 more information on using the new file. Thank you.

**DISCLAIMER:**

Neither the United States Government, nor any agency

thereof, nor any of their contractors, subcontractors or employees make any warranty, expressed or implied,

\* \* \* \* \*  
\* \* \* \* \* W E L C O M E T O T H E  
\* \* \* \* \* U . S . P A T E N T T E X T F I L E  
\* \* \* \* \*  
\* \* \* \* \*  
\* \* \* \* \*  
\* \* \* \* \*

=> s pertussis  
L1 1133 PERTUSSIS  
=> s toxin  
L2 4029 TOXIN  
=> s 11(5n)l2  
L3 226 L1(5A)L2  
=> s mutat? or substitut? or mutan? or mutagen?  
11102 MUTAT?  
402540 SUBSTITUT?  
11912 MUTAN?  
7771 MUTAGEN?  
L4 411817 MUTAT? OR SUBSTITUT? OR MUTAN? OR MUTAGEN?  
=> s repla?c  
'?' TRUNCATION SYMBOL NOT VALID WITHIN 'REPLA?C'  
=> s replac?  
L5 572562 REPLAC?  
=> s s l4 or l5  
L6 832228 L4 OR L5  
75% OF LIMIT FOR TOTAL ANSWERS REACHED  
=> s s l3(5n) l6  
L7 14 L3(5A) L6  
=> d

US PAT NO: 5,705,361 [IMAGE AVAILABLE] L7: 1  
of 14  
DETDESC:  
DETD(10)  
In . . . and biological activities is a protective antigen. Infect. Immun. 58:3337-3347; and Nencioni, L., et al. 1990. Characterization of genetically inactivated \*\*pertussis\*\* \*\*toxin\*\* \*\*mutants\*\* candidates for a new vaccine against whooping cough. Infect. Immun. 58:1308-1315). However, those methods rely on the purification of detoxified.  
. . .  
=> d fro

US PAT NO: 5,705,361 [IMAGE AVAILABLE] L7: 1  
of 14  
DATE ISSUED: Jan. 6, 1998  
TITLE: Hybrid transposon with PT operon, plasmids and bacterial strains therefor  
INVENTOR: Mark Walker, Brunswick, Federal Republic of Germany  
Germany  
ASSIGNEE: Kenneth Timmis, Brunswick, Federal Republic of Germany  
Gesellschaft fur Biotechnologische Forschung mbH GBF, Brunswick, Federal Republic of Germany  
(foreign corp.)  
APPL-NO: 08/137,139  
DATE FILED: Oct. 21, 1993  
PCT-FILED: Apr. 24, 1992  
PCT-NO: PCT/EP92/00910  
371-DATE: Oct. 21, 1993

102 (E) -DATE: Oct. 21, 1993  
PCT-PUB-NO: WO92/19750  
PCT-PUB-DATE: Nov. 12, 1992  
FRN-PRIOR: Federal Republic of Germany41 13 385.4  
Apr. 24, 1991  
INT-CL: [6] C12P 21/02; C12N 1/20; C12N 15/74  
US-CL-ISSUED: 435/69.3, 252.3, 320.1  
US-CL-CURRENT: 435/69.3, 252.3, 320.1  
SEARCH-FLD: 435/320.1, 252.3, 172.3, 69.3; 424/92  
REF-CITED:  
4,883,761 11/1989 U.S. PATENT DOCUMENTS  
Keith et al.  
435/320.1

OTHER PUBLICATIONS  
Walker et al., Infection and Immunity, vol. 59, Nov. 1991, pp. 4238-4248.  
Herrero et al., V. Bacteriol., vol. 172, Nov. 1990, pp. 6557-6567.  
Bloom, Rev. Infect. Diseases, vol. 11, Supplement 2, 1989, pp. 5460-5466.  
Knapp et al., vol. 170, J. Bacteriol., 1988, pp. 5059-5066.  
ART-UNIT: 185  
PRIM-EXMR: James S. Ketter  
LEGAL-REP: Kane, Dalsimer, Sullivan, Kurucz, Levy, Eisele and Richard

ABSTRACT:

The invention relates to a hybrid transposon having a PT operon, to plasmids and to bacterial strains (hosts) therefor.  
12 Claims, 4 Drawing Figures

=> d clm

US PAT NO: 5,705,361 [IMAGE AVAILABLE] L7: 1  
of 14

CLAIMS:

CLMS(1)

We claim:

1. A plasmid for the stable expression of pertussis toxin in a bacterial microorganism which comprises;

a promoterless hybrid transposon wherein a transposase gene is arranged outside of inverted repeats of the transposon; and having a pertussis toxin operon inserted between the inverted repeats;  
said inverted repeats being in shortened form, thereby permitting a further transposition;  
said plasmid permitting a stable insertion of the pertussis toxin operon into a bvg-negative bacterial strain.

CLMS(2)

2. A plasmid according to claim 1 wherein the transposon is a mini-transposon.

CLMS(3)

3. A plasmid according to claim 1 wherein the transposon is from Tn5 and the inverted repeats each have a length of approximately 19 base pairs.

CLMS(4)

4. A plasmid according to claim 1 wherein the operon is selected from the group consisting of a wild-type operon and an operon that encodes a pertussis toxin modification effective against whooping cough.

CLMS(5)

5. A plasmid according to claim 1 wherein the hybrid transposon contains a kanamycin resistance gene.

CLMS(6)

6. A plasmid which is promoterless having the following features and is a suicide vector that has a mini-transposon for pertussis toxin expression for transcriptional fusions with native promoters, having

1. a mob site,
2. an ampicillin resistance gene and
3. in each case, 19 base pairs of the terminal Tn5 end which flank,

4. a kanamycin resistance gene and
5. a promoterless operon; and also having
6. a IS50.sub.R -transposase gene.

#### CLMS (7)

7. A bvg-negative bacterial strain containing a plasmid according to claim 1.

#### CLMS (8)

8. The bacterial strain of claim 7 which is selected from the group consisting of Bordetella pertussis, Bordetella parapertussis and Bordetella bronchiseptica.

#### CLMS (9)

9. A bvg-negative bacterial strain wherein a pertussis toxin operon has been incorporated by means of a plasmid according to one of claims 1, 2 and 3 to 6.

#### CLMS (10)

10. The strain of claim 9 selected from the group of species consisting of Bordetella pertussis, Bordetella parapertussis and Bordetella bronchiseptica.

#### CLMS (11)

11. The strain of claim 10 which is Bordetella bronchiseptica ATCC 10580.

#### CLMS (12)

12. A process for the stable expression of pertussis toxin which comprises; culturing a bvg-negative bacterial strain containing a plasmid according to one of claims 1, 2 and 3 to 6.

=> s (arg or arginine) (w)9 or arg9 or arginine9

8246 ARG  
13998 ARGinine  
1660508 9

49 (ARG OR ARGinine) (W) 9  
8 ARG9

1 ARGinine9

L8 57 (ARG OR ARGinine) (W) 9 OR ARG9 OR ARGinine9

=> s 18(p)11

L9 1 L8(P)L1

=> s 13 and 18

L10 1 L3 AND L8

=> s vaccine

L11 4528 VACCINE

=> s 111(p)13

L12 48 L11(P)L3

=> d 1-3

1. 5,725,860, Mar. 10, 1998, Method for reducing the risk of developing diabetes; Robert Bartlett Elliott, 424/240.1; 514/2, 21 [IMAGE AVAILABLE]

2. 5,705,361, Jan. 6, 1998, Hybrid transposon with PT operon, plasmids and bacterial strains therefor; Mark Walker, et al., 435/69.3, 252.3, 320.1 [IMAGE AVAILABLE]

3. 5,695,766, Dec. 9, 1997, Highly virulent porcine reproductive and respiratory syndrome viruses which produce lesions in pigs and vaccines that protect pigs against said syndrome; Prem S. Paul, et al., 424/204.1, 209.1, 218.1, 815; 435/235.1, 236, 237, 238, 239 [IMAGE AVAILABLE]

=> d kwic

US PAT NO:  
of 48

5,725,860 [IMAGE AVAILABLE]

L12: 1

DETD(45)

DETD(45)

The conclusions we draw from this are that the pertussis  
\*\*vaccine\*\* is  
the active adjuvant in the mixed, triple \*\*vaccine\*\*, and that  
the  
diphtheria and tetanus \*\*vaccine\*\* are without effect. It also  
appears  
that A-chain itself, or in combination with diphtheria and/or  
tetanus  
\*\*vaccine\*\* is ineffective. The pertussis \*\*vaccine\*\* used was  
a  
"cellular" \*\*vaccine\*\* containing many defined, and some  
undefined  
components, and so we are unable to specify which component or  
components  
of the \*\*vaccine\*\* contain(s) the active principle. Indeed,  
diphtheria  
and/or tetanus vaccines may be required in addition as "co-  
factors". It  
may be relevant that for a long time \*\*pertussis\*\* \*\*toxin\*\*  
has been  
known to stimulate insulin secretion from the islets of  
Langerhans,  
probably through the "G protein" control sequence.

=> d 2-3 kwic

US PAT NO:  
of 48

5,705,361 [IMAGE AVAILABLE]

L12: 2

SUMMARY:

BSUM(3)

The invention relates to a plasmid for bacterial expression  
of  
\*\*pertussis\*\* \*\*toxin\*\*, useful in \*\*vaccine\*\* preparation.

SUMMARY:

BSUM(5)

Increasing . . . has in turn resulted in an increase in  
the incidence  
of the disease. It is imperative that an effective,  
nonreactogenic  
\*\*vaccine\*\* of high acceptability be developed. Several  
potentially  
protective antigens are candidates for inclusion in a purified  
multi-component \*\*vaccine\*\*, including toxins, such as  
\*\*pertussis\*\*  
\*\*toxin\*\* (PT) and adenylate cyclase, and also several cell  
surface or  
secreted antigens like filamentous haemagglutinin and  
serotype-specific  
fimbriae (Robinson, A., et al. 1985. Pertussis \*\*vaccine\*\*:  
status and future prospects. \*\*Vaccine\*\* 3:11-22; and  
Robinson. A., and  
I. A. E. Ashworth. 1988. Acellular and defined-component  
vaccines against  
pertussis, p. 399-417. In A. . . is hampered by antigenic  
variation  
controlled by the bvg-positive regulatory locus (Gross, R.,  
and R.  
Rappuoli. 1988. Positive regulation of \*\*pertussis\*\* \*\*toxin\*\*  
expression. Proc. Natl. Acad. Sci. 85:3913-3917; Knapp, S.,  
and J. J.  
Mekalanos. 1988. Two trans-acting regulatory genes (vir and  
mod). . .  
E. coli under the control of the lambda P.sub.L promoter  
(Burnette, W.  
N., et al. 1988. Direct expression of Bordetella \*\*pertussis\*\*  
\*\*toxin\*\*  
subunits to high levels in Escherichia coli. Biotechnol.  
6:699-706). The  
serotype 2 fimbrial sub-unit has also been expressed with the.  
. .  
also extremely unstable, being subject to both plasmid loss  
and gene  
deletions (Locht, C., et al. 1986. Molecular cloning of  
\*\*pertussis\*\*  
\*\*toxin\*\* genes. Nucl. Acids Res. 14:3251-3261).

DETD(10)

DETD(10)

In . . . that methods for the genetic detoxification of PT  
have  
recently been developed by several groups (Kimura, A., et al.  
1990.



**\*\*Pertussis\*\*** \*\*toxin\*\* analog with reduced enzymatic and biological activities is a protective antigen. Infect. Immun. 58:3337-3347; and Nencioni, L., et al. 1990. Characterization of genetically inactivated **\*\*pertussis\*\*** \*\*toxin\*\* mutants: candidates for a new **\*\*vaccine\*\*** against whooping cough. Infect. Immun. 58:1308-1315). However, those methods rely on the purification of detoxified PT from B. pertussis. The . . of bvg-negative strains of B. bronchiseptica in combination with methods for the genetic detoxification of PT will lead to a **\*\*vaccine\*\*** component of high purity and low reactogenicity.

US PAT NO: 5,695,766 [IMAGE AVAILABLE] L12: 3 of 48

DETDESC:

DETD(51)

The composition containing the present **\*\*vaccine\*\*** may be administered in conjunction with an adjuvant. An adjuvant is a substance that increases the immunological response to the present **\*\*vaccine\*\*** when combined therewith. The adjuvant may be administered at the same time and at the same site as the **\*\*vaccine\*\*** or at a different time, for example, as a booster. Adjuvants also may advantageously be administered to the animal in a manner or at a site or location different from the manner, site or location in which the **\*\*vaccine\*\*** is administered. Adjuvants include aluminum hydroxide, aluminum potassium sulfate, heat-stable enterotoxin isolated from Escherichia coli, cholera toxin or the B subunit thereof, diphtheria **\*\*toxin\*\***, tetanus **\*\*toxin\*\***, **\*\*pertussis\*\*** **\*\*toxin\*\***, Freund's incomplete adjuvant, Freund's complete

adjuvant, and the like. Toxin-based adjuvants, such as diphtheria **\*\*toxin\*\***, tetanus **\*\*toxin\*\*** and **\*\*pertussis\*\*** **\*\*toxin\*\***, may be inactivated prior to use, for example, by treatment with formaldehyde.

=> s (514/12 or 424/240.1 or 424/190.1 or 424/94.5)/ccls

1716 514/12/CCLS  
45 424/240.1/CCLS  
80 424/190.1/CCLS

61 424/94.5/CCLS

L13 1882 (514/12 OR 424/240.1 OR 424/190.1 OR 424/94.5)/CCLS

=> s l3 and l13

L14 47 L3 AND L13

=> d 1-3

1. 5,736,143, Apr. 7, 1998, Anti-inflammatory, tolerogenic and immunoinhibiting properties of carbohydrate binding-peptides; Louis D. Heerze, et al., \*\*424/190.1\*\*\*, 275.1; 514/885 [IMAGE AVAILABLE]

2. 5,731,151, Mar. 24, 1998, Regulator of contact-mediated hemolysin; C. Harold King, et al., 435/6; \*\*424/190.1\*\*\*, 200.1, 248.1; 435/41, 69.1, 71.1; 536/25.3 [IMAGE AVAILABLE]

3. 5,725,860, Mar. 10, 1998, Method for reducing the risk developing diabetes; Robert Bartlett Elliott, \*\*424/240.1\*\*\*, 514/2, 21 [IMAGE AVAILABLE]

=> s (toxin/clm) or holotoxin/clm or s1/clm

724 TOXIN/CLM  
9 HOLOTOXIN/CLM  
1280 S1/CLM

L15 1997 (TOXIN/CLM) OR HOLOTOXIN/CLM OR S1/CLM

=> s l14 and l15

L16 29 L14 AND L15

=> d his

(FILE 'USPAT' ENTERED AT 13:58:18 ON 04 MAY 1998)

L1 1133 S PERTUSSIS  
L2 4029 S TOXIN  
L3 226 S L1(5N)L2  
L4 411817 S MUTAT? OR SUBSTITUT? OR MUTAN? OR MUTAGEN?  
L5 572562 S REPLAC?  
L6 832228 S L4 OR L5  
L7 14 S L3(5N) L6  
L8 57 S (ARG OR ARGinine) (W) 9 OR ARG9 OR ARGinine9  
L9 1 S L8(P)L1  
L10 1 S L3 AND L8  
L11 4528 S VACCINE  
L12 48 S L11(P)L3  
L13 1882 S (514/12 OR 424/240.1 OR 424/190.1 OR  
424/94.5)/CCLS  
L14 47 S L3 AND L13  
L15 1997 S (TOXIN/CLM) OR HOLOTOXIN/CLM OR S1/CLM  
L16 29 S L14 AND L15

=> logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF  
LOGOFF? (Y)/N/HOLD:Y

U.S. Patent & Trademark Office LOGOFF AT 14:04:47 ON 04 MAY  
1998